



Review Article

Effects of Herbal Medicine in the Treatment of Poultry Coccidiosis

Mohammad Reza Eftekhari Hasan Abad , and Abolfazl Ghaniei* 

Department of Clinical Sciences, Faculty of Veterinary Medicine, Ferdowsi University of Mashhad, Mashhad, Iran

* **Corresponding author:** Abolfazl Ghaniei, Department of Clinical Sciences, Faculty of Veterinary Medicine, Ferdowsi University of Mashhad, Mashhad, Iran.
Email: ghaniei@um.ac.ir

ARTICLE INFO

Article History:

Received: 26/01/2023

Accepted: 02/03/2023



Keywords:

Coccidiosis

Eimeria

Herbal medicine

Poultry

ABSTRACT

Poultry coccidiosis is a common and economically significant disease that affects poultry worldwide. Current treatment options for coccidiosis include using chemical agents such as ionophores and antiparasitic such as Metronidazole. However, there is growing interest in alternative treatments, such as herbal medicines, due to concerns about antiparasitic resistance and residues in poultry products. This review article aims to provide an overview of herbal medicines as potential treatments for poultry coccidiosis. The article summarizes the current research on herbal medicines for coccidiosis and highlights the potential benefits and limitations of using herbal medicines as an alternative to conventional treatments. The article explores the effectiveness of various herbal medicines in treating poultry coccidiosis, explaining how they work and presenting research findings on their efficacy. It also addresses challenges related to using herbal medicines, such as establishing standardized doses and conducting clinical trials. The review concludes by summarizing the main points and recommending further research on herbal medicines for poultry coccidiosis. Ultimately, the review underscores the potential of herbal medicines as alternative treatments and emphasizes the importance of continued investigation in this field.

1. Introduction

Poultry coccidiosis is a widespread disease affecting domestic and wild birds worldwide¹. The disease is caused by different species of *Eimeria* (protozoa) that infect the intestinal tract of chickens, turkeys, ducks, and other poultry species². The infection can lead to severe damage to the intestinal lining, resulting in malabsorption of nutrients, diarrhea, and dehydration. In addition, coccidiosis in poultry results in reduced weight gain, decreased egg production, and increased mortality rates, resulting in significant economic losses to the poultry industry³. The most common approach to control coccidiosis in poultry is using chemical agents, including ionophores and antiparasitic, such as Metronidazole⁴. However, the overuse of antibiotics in animal production has been linked to the development of antibiotic-resistant bacteria, which can pose a threat to public health⁵. In addition, antibiotic residues in poultry products have raised concerns among consumers, leading to a shift towards using natural and organic products in animal

production⁶.

Herbal medicines have been used as alternative treatments for various diseases for centuries^{7,8}. Plants, such as *Echinacea purpurea* and *Glycyrrhiza glabra* contain a diverse range of bioactive compounds, including alkaloids, flavonoids, and terpenoids, which have a wide range of pharmacological activities⁹⁻¹¹. Herbal medicines are known to have fewer side effects than synthetic drugs, making them a potential alternative to chemical agents for controlling poultry coccidiosis. In recent years, there has been a growing interest in using herbal medicines to control poultry coccidiosis, with several studies investigating the potential of herbs and plant extracts as alternative treatments^{12,13}. Multiple studies have demonstrated that specific herbs and plant extracts possess properties that can combat coccidiosis in poultry. These findings have highlighted the potential of using these herbs as alternative treatments for the disease^{14,15}.

Herbs with anti-coccidial properties include turmeric (*Curcuma longa*), neem (*Azadirachta indica*), oregano (*Origanum vulgare*), garlic (*Allium sativum*), sweet wormwood (*Artemisia annua*), and thyme (*Thymus vulgaris*)¹⁶⁻¹⁸. These herbs have been shown to possess antimicrobial, antioxidant, and anti-inflammatory properties and effectively reduce the severity of coccidiosis in poultry¹⁹. The use of herbal medicines as alternative treatments for poultry coccidiosis has the potential to reduce the reliance on chemical agents, mitigate the development of antiparasitic resistance, and promote the use of natural and organic products in animal production. However, further research is needed to determine these herbal medicines' optimal dosages, safety, and efficacy in controlling poultry coccidiosis.

The purpose of this review was to offer a thorough and detailed summary of the existing research on herbal medicines and their potential as treatments for poultry coccidiosis. This review aimed to inform and guide future research on using herbal medicines as alternative treatments for poultry coccidiosis by summarizing the existing knowledge on this topic.

2. Herbal medicines history

Herbal medicines, such as *Echinacea purpurea* and *Glycyrrhiza glabra* have been used in traditional medicine systems for centuries to treat various diseases, including infections^{20,21}. There has been growing interest in using herbal medicines as alternative treatments for poultry coccidiosis in recent years^{22,23}. Herbal medicines are plant-derived products that contain active compounds, such as curcumin and allicin with therapeutic properties. These compounds can act on different targets in the body to exert their beneficial effects, including anti-inflammatory, antimicrobial, and antioxidant properties²⁴. Herbal medicines, with their longstanding use in traditional medicine systems like Chinese medicine. In the context of poultry diseases, including poultry coccidiosis, there is increasing interest in investigating the efficacy of herbal medicines as alternative treatments²⁵.

In modern times, herbal medicines have gained attention as alternative treatments for animal diseases due to concerns about developing antiparasitic resistance and using chemical agents in animal production²⁶.

Research on using herbal medicines for poultry coccidiosis is still in its early stages, but several studies have shown promising results^{27,28}. These studies have investigated the potential of different herbs and plant extracts for controlling the disease and reducing its impact on poultry production.

3. Herbal medicines with efficacy against poultry coccidiosis

Several herbal medicines are effective in treating poultry coccidiosis. These herbs contain active compounds that have anti-coccidial properties and can act on different targets in the body to reduce the severity of the disease.

3.1. Garlic (*Allium sativum*)

Garlic is a popular herb used for culinary and medicinal purposes for thousands of years. It is a member of the *Allium* genus, including onions, leeks, and chives²⁹. Garlic is known for its distinct aroma and flavor from sulfur-containing compounds such as allicin^{30,31}. Allicin, the active compound in garlic, effectively inhibits the growth and development of *Eimeria* parasites^{32,33}. It achieves this by disrupting the parasite's cell membrane and inhibiting its ability to replicate³⁴. In addition, allicin has been shown to stimulate the immune system, which can help chickens fight against parasitic infections³⁵.

3.2. Thyme

Thyme is a culinary herb used for centuries for its distinct flavor and aroma. It belongs to the mint family, *Lamiaceae*, and is commonly found in Mediterranean and Middle Eastern cuisine³⁶. Thyme has also been used for medicinal purposes, with its essential oil being used as an antiseptic and disinfectant both in humans and veterinary medicine^{37,38}.

Recent studies have shown that thyme possesses anti-coccidial properties and can effectively control poultry coccidiosis^{39,40}. A study indicated that an active compound in thyme (thymol) is particularly effective in reducing the severity of *Eimeria* infections in chickens⁴¹. Thymol is a natural phenol that has been found to have antimicrobial, antifungal, and antiparasitic properties⁴² by disrupting the life cycle of *Eimeria* parasites⁴³. It interferes with the parasite's ability to reproduce and survive in the chicken's intestine, thus reducing the severity of the infection⁴⁴. Thymol has also been found to have immunomodulatory effects, stimulating the chicken's immune system to fight off the infection⁴⁵.

In addition to thymol, thyme contains other bioactive compounds that may contribute to its anti-coccidial properties⁴⁶. These include carvacrol, rosmarinic acid, and flavonoids, which have been found to have antimicrobial and anti-inflammatory properties⁴⁷.

3.3. Oregano

It has recently gained attention for its potential use in treating poultry coccidiosis^{48,49}. The active compounds in oregano, carvacrol, and thymol, have been shown to possess potent anti-coccidial properties^{50,51} by disrupting the development of the *Eimeria* parasite, which causes coccidiosis in chickens⁵².

Several studies have shown that feeding oregano to chickens infected with *Eimeria* can reduce the severity of the disease and improve their overall health^{53,54}. In one study, researchers found that supplementing infected chickens' diet with oregano essential oil reduced the number of the *Eimeria* spp. oocysts shed in their feces⁵⁵. The treated chickens also had higher weight gains than infected chickens that did not receive oregano.

Another study tested the efficacy of oregano extract in treating poultry coccidiosis⁵⁶. The results indicated that oregano extract significantly reduced the severity of the disease and the number of oocysts shed in the feces of infected chickens. The researchers concluded that oregano extract could be a safe and effective alternative treatment for poultry coccidiosis⁵⁷.

3.4. Neem

Neem, also known as Indian lilac, is a tree that is widely used in traditional medicine in India for its therapeutic properties⁵⁸. Its active compounds, azadirachtin, and nimbin, have been found to have anti-coccidial properties, making it a promising candidate for the treatment of poultry coccidiosis⁵⁹.

A study on the efficacy of neem against poultry coccidiosis found that feeding neem to infected chickens reduced the severity of the coccidiosis disease and improved their weight gain compared to infected chickens that did not receive neem⁶⁰. The active compounds found in neem were identified as responsible for their anti-coccidial properties, as they were observed to inhibit the growth of *Eimeria* parasites in chickens.

Neem has also been found to have other health benefits for chickens^{61,62}. It can act as an immune booster and promote gut health, which is crucial for preventing poultry diseases, including coccidiosis^{63,64}.

In addition to its anti-coccidial properties, neem has also been found to possess other medicinal properties⁶⁵. It has anti-inflammatory, antioxidant, and antibacterial properties, making it useful for treating various diseases in chickens⁶⁶.

3.5. Curcumin

Curcumin, the active compound in turmeric, has also been studied for its potential anti-coccidial properties in chickens⁶⁷. Turmeric is a spice commonly used in Indian cuisine and has been used for its medicinal properties in traditional medicine systems⁶⁸.

A study on the efficacy of curcumin against poultry coccidiosis found that feeding curcumin to infected chickens reduced the number of *Eimeria* oocysts shed in their feces and improved their weight gain compared to infected chickens not receiving curcumin⁶⁹. The researchers attributed the anti-coccidial properties of curcumin to its ability to modulate the host immune response and inhibit the development of *Eimeria* parasites in chickens.

Curcumin has also been found to have other health benefits for chickens⁷⁰. It can act as an antioxidant, anti-inflammatory, and antibacterial agent, which makes it useful for the prevention and treatment of various poultry diseases, including coccidiosis⁷¹.

Moreover, curcumin is a natural and organic product safe for chickens and humans⁷². Its safety and efficacy make it a valuable natural and organic treatment for poultry diseases.

3.6. *Artemisia annua*

Artemisia annua, commonly known as sweet wormwood. This herb has been used in traditional Chinese medicine for centuries and has been shown to possess anti-coccidial properties⁷³. The active ingredient in *Artemisia annua* is artemisinin, which has been shown to have anti-coccidial activity against different species of *Eimeria* species⁷⁴. Another study aimed to compare the effects of different *Artemisia annua* extracts on the sporulation rate of mixed oocysts of *Eimeria acervulina*, *Eimeria necatrix*, and *Eimeria tenella*⁷⁵. Three types of *A. annua* extracts (petroleum ether, ethanol, and water) were prepared and analyzed. The results showed that petroleum ether and ethanol extracts inhibited sporulation at certain concentrations.

4. Mechanisms of action

Herbal medicines have been shown to possess a range of mechanisms of action that contribute to their efficacy against poultry coccidiosis. For example, many herbs contain active compounds with anti-inflammatory properties, such as curcumin in turmeric and gingerols in ginger⁷⁶. Inflammation plays a critical role in the immune response to coccidiosis. It serves as a key component in the body's defense mechanism against the infection caused by coccidia parasites. By reducing inflammation, these herbs can help alleviate the symptoms of the disease and support the chicken's overall health.

Other herbal medicines have anti-parasitic properties, which means they can directly kill or inhibit the growth of coccidian parasites. For example, thymol and carvacrol in oregano have been shown to have strong anti-parasitic activity, while allicin in garlic has been found to inhibit the growth of *Eimeria*⁷⁷. These anti-parasitic properties can help reduce the number of parasites in the chicken's gut, reducing the severity of the disease and supporting the chicken's recovery.

Finally, some herbs have immune-modulating properties, which can help regulate the chicken's immune response to the parasite. Neem has been shown to have the ability to enhance the production of specific cytokines, which are important signaling molecules involved in regulating the immune response. This suggests that neem may contribute to strengthening the immune response against coccidiosis and other infections⁷⁸.

By modulating the immune response, these herbs can help the chicken fight against the infection more effectively and reduce the damage caused by the parasite⁷⁹.

Overall, the range of mechanisms by which herbal medicines act against poultry coccidiosis highlights the potential benefits of using these natural remedies in place of or in conjunction with chemical treatments.

5. Future and Challenges

The use of herbal medicines for controlling poultry coccidiosis has shown promising results, but this approach

has several challenges and limitations. One of the main challenges is the need for standardized extraction methods and dosing protocols for herbal medicines⁸⁰. This makes it difficult to compare results across different studies and hinders the development of consistent and effective treatments. Additionally, the variability in the composition of herbal extracts due to differences in soil, climate, and harvesting methods can further complicate the standardization process⁸¹.

Another challenge is the need for more research to establish the safety and efficacy of herbal medicines⁸². Although many studies have reported positive results, there still needs to be comprehensive toxicity studies and clinical trials that can provide more robust evidence of their safety and effectiveness⁸³. There needs to be more funding for research on herbal medicines to progress in this area.

The regulatory environment also challenges the use of herbal medicines in animal production. The approval process for herbal medicines is often more complex and time-consuming than chemical agents⁸⁴. Additionally, the lack of standardized regulations across different countries can create barriers to the international trade of herbal medicines⁸⁵.

Moreover, there is a need for more awareness and education among farmers and veterinarians about using herbal medicines to control poultry coccidiosis⁸⁶. Many farmers may hesitate to adopt alternative treatments due to a lack of familiarity and understanding of their mechanisms of action and potential benefits.

Despite these challenges, the use of herbal medicines for controlling poultry coccidiosis has the potential to offer several benefits over chemical agents. Herbal medicines are generally considered safe and have a lower risk of developing antiparasitic resistance than chemical agents. They are also compatible with organic and natural production systems, which are becoming increasingly popular among consumers. Moreover, using herbal medicines can help reduce the environmental impact of animal production by reducing the amount of chemical residues in animal products⁸⁷.

6. Conclusion

In conclusion, herbal medicines have shown great potential as alternative treatments for poultry coccidiosis. Various herbs and plant extracts have been shown to possess anti-coccidial properties. They can be used to reduce the reliance on chemical agents, mitigate the development of antiparasitic resistance, and promote the use of natural and organic products in animal production. Garlic, thyme, oregano, and neem are herbal medicines effective in treating poultry coccidiosis. The mechanisms by which herbal medicines act against poultry coccidiosis include their anti-inflammatory, anti-parasitic, and immune-modulating properties. However, further research is needed to better understand the mechanisms of action and optimize the use of these herbal medicines to control poultry coccidiosis.

Future challenges in using herbal medicines for poultry

coccidiosis include standardization of products, development of effective delivery methods, and identification of optimal dosages. Additionally, regulatory issues, such as lack of approval for herbal medicines in some countries, may limit their widespread usage. In conclusion, while there are several challenges and limitations to using herbal medicines for controlling poultry coccidiosis, their potential benefits make them a promising area for further research and development. It is important to continue to invest in research to establish the safety and efficacy of herbal medicines, standardize extraction methods and dosing protocols, and raise awareness and education among farmers and veterinarians about their potential benefits.

Declarations

Competing interests

There is no conflict of interest.

Authors' contribution

The final manuscript draft was reviewed by all authors, who also gave their approval.

Funding

No funding.

Ethical considerations

Ethical issues (including plagiarism, consent to publish, misconduct, data fabrication and/or falsification, double publication and/or submission, and redundancy) have been checked by all the authors.

Acknowledgments

The authors wish to acknowledge everyone who helped during the writing of this manuscript.

References

1. Lozano J, Anaya A, Palomero Salinero A, Lux Hoppe EG, Gomes L, Paz-Silva A, et al. Gastrointestinal parasites of free-range chickens: A worldwide issue. *Bulletin of University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca. Vet Med.* 2019; 76(2): 110-117. Available at: <https://www.repository.utl.pt/handle/10400.5/21753>
2. McDougald LR, Cervantes HM, Jenkins MC, Hess M, and Beckstead R. Protozoal infections. In: Swayne DE, Boulianne M, Logue CM, McDougald LR, Nair V, Suarez DL, et al., editors. *Diseases of poultry*. 2020 Nov; p. 1192-1254. DOI: [10.1002/9781119371199.ch28](https://doi.org/10.1002/9781119371199.ch28)
3. Blake DP, Knox J, Dehaeck B, Huntington B, Rathinam T, Ravipati V, et al. Re-calculating the cost of coccidiosis in chickens. *Vet Res.* 2020 Nov; 51: 115. Available at: <https://link.springer.com/article/10.1186/s13567-020-00837-2>
4. Chapman H, Jeffers T, and Williams R. Forty years of monensin for the control of coccidiosis in poultry. *Poult Sci.* 2010 Sep; 89(9): 1788-

1801. DOI: [10.3382/ps.2010-00931](https://doi.org/10.3382/ps.2010-00931)
5. Mirzaei A, Razavi SA, Babazadeh D, Laven R, and Saeed M. Roles of probiotics in farm animals: A review. *Farm Anim Health Nut*. 2022 Sep; 1(1): 17-25. Available at: <https://fahn.rovedar.com/index.php/FAHN/article/view/8>
6. Walsh C, and Fanning S. Antimicrobial resistance in foodborne pathogens-A cause for concern?. *Curr Drug Targets*. 2008; 9(9): 808-815. DOI: [10.2174/138945008785747761](https://doi.org/10.2174/138945008785747761)
7. Sadr S, Ahmadi PS, Kasaei M, Landi MG, Borji H, and Adhami G. Potential of anthelmintic herbal drugs against gastrointestinal nematodes in farm animals: A review. *Farm Anim Health Nut*. 2022 Sep; 1(1): 26-30. Available at: <https://fahn.rovedar.com/index.php/FAHN/article/view/9>
8. Sadr S, Gharib A, Lotfalizadeh N, Hajjafari A, Ahmadi PS, and Borji H. Phytosomes: A promising nanocarrier for enhanced delivery of herbal compounds in cancer therapy. *J Lab Anim Res*. 2022 Dec; 1(1): 26-32. Available at: <https://jlar.rovedar.com/index.php/JLAR/article/view/8>
9. Ghafouri SA, Ghaniei A, Tamannaie AET, Sadr S, Charbgo A, Ghiassi S, et al. Evaluation of therapeutic effects of an herbal mixture (*Echinacea purpurea* and *Glycyrrhiza glabra*) for treatment of clinical coccidiosis in broilers. *Vet Med Sci*. 2023 Mar; 9(2): 829-836. DOI: [10.1002/2Fvms.3.971](https://doi.org/10.1002/2Fvms.3.971)
10. Ghaniei A, Ghafouri SA, Sadr S, and Hassanbeigi N. Investigating the preventive effect of herbal medicine (*Allium sativum*, *Artemisia annua*, and *Quercus infectoria*) against coccidiosis in broiler chickens. *J World's Poult Res*. 2023 Mar; 13(1): 96-102. DOI: [10.36380/jwpr.2023.10](https://doi.org/10.36380/jwpr.2023.10)
11. Sadr S, Ghafouri SA, Ghaniei A, Jami Moharreri D, Zeinali M, Qaemifar N, et al. Treatment of avian trichomoniasis by tannin-based herbal mixture (*Artemisia annua*, *Quercus infectoria*, and *Allium Sativum*). *J World's Poult Sci*. 2022; 1(2): 32-39. Available at: <https://jwps.rovedar.com/index.php/JWPS/article/view/6>
12. Muthamilselvan T, Kuo TF, Wu YC, and Yang WC. Herbal remedies for coccidiosis control: A review of plants, compounds, and anticoccidial actions. *Evid-based Complement Altern Med: eCAM*. 2016 Jun; 2016: 2657981. DOI: [10.1155/2016/2657981](https://doi.org/10.1155/2016/2657981)
13. Hartady T, Syamsunarno MRA, Priosoeryanto BP, Jasni S, and Balia RL. Review of herbal medicine works in the avian species. *Vet world*. 2021 Nov; 14(11): 2889. DOI: [10.14202/vetworld.2021.2889-2906](https://doi.org/10.14202/vetworld.2021.2889-2906)
14. Jamil M, Aleem MT, Shaikat A, Khan A, Mohsin M, Rehman TU, et al. Medicinal plants as an alternative to control poultry parasitic diseases. *Life*. 2022 Mar; 12(3): 449. DOI: [10.3390/life12030449](https://doi.org/10.3390/life12030449)
15. Madlala T, Okpeku M, and Adeleke MA. Understanding the interactions between *Eimeria* infection and gut microbiota, towards the control of chicken coccidiosis: A review. *Parasite*. 2021 Jun; 28: 48. DOI: [10.1051/2Fparasite/2F2021047](https://doi.org/10.1051/2Fparasite/2F2021047)
16. Pop LM, Varga E, Coroian M, Nedişan ME, Mircean V, Dumitrache MO, et al. Efficacy of a commercial herbal formula in chicken experimental coccidiosis. *Parasit vectors*. 2019; 12: 243. DOI: [10.1186/s13071-019-3595-4](https://doi.org/10.1186/s13071-019-3595-4)
17. Christaki E, Bonos E, Giannenas I, and Florou-Paneri P. Aromatic plants as a source of bioactive compounds. *Agriculture*. 2012 Sep; 2(3): 228-243. DOI: [10.3390/agriculture2030228](https://doi.org/10.3390/agriculture2030228)
18. Anthony JP, Fyfe L, and Smith H. Plant active components—a resource for antiparasitic agents? *Trends Parasitol*. 2005 Oct; 21(10): 462-468. DOI: [10.1016/j.pt.2005.08.004](https://doi.org/10.1016/j.pt.2005.08.004)
19. Adaszyńska-Skwirzyńska M, and Szczerbińska D. Use of essential oils in broiler chicken production—a review. *Ann Anim Sci*. 2017 May; 17(2): 317. DOI: [10.1515/aoas-2016-0046](https://doi.org/10.1515/aoas-2016-0046)
20. Ahmad Khan MS, and Ahmad I. Herbal medicine: Current trends and future prospects. *New Look to Phytomedicine*, Chapter 1. 2019. p. 3-13. DOI: [10.1016/B978-0-12-814619-4.00001-X](https://doi.org/10.1016/B978-0-12-814619-4.00001-X)
21. Singh R. Medicinal plants: A review. *J Plant Sci*. 2015 Jan; 3(1-1): 50-55. Available at: <https://article.sciencepublishinggroup.com/html/10.11648/jpps.2015030101.18.html>
22. Abebe E, and Gugs G. A review on poultry coccidiosis. *Abyssinia J Sci Technol*. 2018 May; 3(1): 1-12. Available at: <https://abjol.org.et/index.php/ajst/article/view/76>
23. Quiroz-Castañeda RE, and Dantán-González E. Control of avian coccidiosis: Future and present natural alternatives. *Biomed Res Int*. 2015 Feb; 2015: 430610. DOI: [10.1155/2015/430610](https://doi.org/10.1155/2015/430610)
24. Quiroz-Castañeda RE. Avian coccidiosis, new strategies of treatment. *Farm animals diseases, recent omic trends and new strategies of treatment*. 2018 Mar. p. 119. Available at: <https://www.intechopen.com/books/5543>
25. Giannenas I, Sidiropoulou E, Bonos E, Christaki E, and Florou-Paneri P. The history of herbs, medicinal and aromatic plants, and their extracts: Past, current situation and future perspectives. *Feed additives*, Chapter 1. 2020. p. 1-18. DOI: [10.1016/B978-0-12-814700-9.00001-7](https://doi.org/10.1016/B978-0-12-814700-9.00001-7)
26. Cragg GM, and Newman DJ. Natural products: A continuing source of novel drug leads. *Biochim Biophys Acta Gen Subj*. 2013 Jun; 1830(6): 3670-3695. DOI: [10.1016/j.bbagen.2013.02.008](https://doi.org/10.1016/j.bbagen.2013.02.008)
27. Naidoo V, McGaw LJ, Bisschop S, Duncan N, and Eloff JN. The value of plant extracts with antioxidant activity in attenuating coccidiosis in broiler chickens. *Vet Parasitol*. 2008 May; 153(3-4): 214-219. DOI: [10.1016/j.vetpar.2008.02.013](https://doi.org/10.1016/j.vetpar.2008.02.013)
28. Shirley M, and Lillehoj H. The long view: A selective review of 40 years of coccidiosis research. *Avian Pathol*. 2012 Apr; 41(2): 111-121. DOI: [10.1080/03079457.2012.666338](https://doi.org/10.1080/03079457.2012.666338)
29. Sharifi-Rad J, Mnayer D, Tabanelli G, Stojanović-Radić Z, Sharifi-Rad M, Yousaf Z, et al. Plants of the genus *Allium* as antibacterial agents: From tradition to pharmacy. *Cell Mol Biol*. 2016 Aug; 62(9): 57-68. Available at: <https://pubmed.ncbi.nlm.nih.gov/27585263/>
30. Bastaki SM, Ojha S, Kalasz H, and Adeghate E. Chemical constituents and medicinal properties of *Allium* species. *Molec cell biochem*. 2021 Aug; 476(12): 4301-4321 DOI: [10.1007/s11010-021-04213-2](https://doi.org/10.1007/s11010-021-04213-2)
31. Lengbiye EM, Mbadiko CM, Falanga CM, Matondo A, Inkoto CL, Ngoyi EM, et al. Antiviral activity, phytochemistry and toxicology of some medically interesting *Allium* species: A mini review. *Int J Pathogen Res*. 2020 Dec; 5(4): 64-77. DOI: [10.9734/ijpr/2020/v5i430145](https://doi.org/10.9734/ijpr/2020/v5i430145)
32. Abd-ELrahman SM, Mohamed SAA, Mohamed SE, El-Khadragy MF, Dyab AK, Hamad N, et al. Comparative effect of allicin and alcoholic garlic extract on the morphology and infectivity of *Eimeria tenella* oocysts in chickens. *Animals*. 2022 Nov; 12(22): 3185. DOI: [10.3390/ani12223185](https://doi.org/10.3390/ani12223185)
33. Bauri R, Tigga MN, and Kullu SS. A review on use of medicinal plants to control parasites. *Indian J Nat Prod Resour*. [Formerly Natural Product Radiance (NPR)]. 2015 Dec; 6(4): 268-277. DOI: [index.php/IJNPR/article/view/8837](https://doi.org/10.1007/978-93-8043-883-7)
34. El-Khtam A, Abd El Latif A, and El-Hewaity M. Efficacy of turmeric (*Curcuma longa*) and garlic (*Allium sativum*) on *Eimeria* species in broilers. *Int J Basic Appl Sci*. 2014 Oct; 3(3): 349. DOI: [10.14419/ijbas.v3i3.3142](https://doi.org/10.14419/ijbas.v3i3.3142)
35. Arreola R, Quintero-Fabián S, López-Roa RI, Flores-Gutiérrez EO, Reyes-Grajeda JP, Carrera-Quintanar L, et al. Immunomodulation and anti-inflammatory effects of garlic compounds. *J Immunol Res*. 2015 Apr; 2015: 401630. DOI: [10.1155/2015/401630](https://doi.org/10.1155/2015/401630)
36. Kowalczyk A, Przygodna M, Sopata S, Bodalska A, and Fecka I. Thymol and thyme essential oil—new insights into selected therapeutic applications. *Molecules*. 2020 Sep; 25(18): 4125. DOI: [10.3390/molecules25184125](https://doi.org/10.3390/molecules25184125)
37. Dauqan EM, and Abdullah A. Medicinal and functional values of thyme (*Thymus vulgaris* L.) herb. *J Appl Biol Biotechnol*. 2017 Mar-Apr; 5(2): 17-22. DOI: [10.7324/JABB.2017.50203](https://doi.org/10.7324/JABB.2017.50203)
38. Hosseinzadeh S, Jafarikukhdan A, Hosseini A, and Armand R. The application of medicinal plants in traditional and modern medicine: A review of *Thymus vulgaris*. *Int J Clin Med*. 2015 Sep; 6(09): 635-642. DOI: [10.4236/ijcm.2015.69084](https://doi.org/10.4236/ijcm.2015.69084)
39. Bozkurt M, Giannenas I, Küçükyılmaz K, Christaki E, and Florou-Paneri P. An update on approaches to controlling coccidia in poultry using botanical extracts. *Br Poult Sci*. 2013 Dec; 54(6): 713-727. DOI: [10.1080/00071668.2013.849795](https://doi.org/10.1080/00071668.2013.849795)
40. Idris M, Abbas R, Masood S, Rehman T, Farooq U, Babar W, et al. The potential of antioxidant rich essential oils against avian coccidiosis. *World's Poult Sci J*. 2017 Mar; 73(1): 89-104. DOI: [10.1017/S0043933916000787](https://doi.org/10.1017/S0043933916000787)
41. Jitviriyanon S, Phanthong P, Lomarar P, Bunyapraphatsara N, Porntrakulpipat S, and Paraksa N. *In vitro* study of anti-coccidial activity of essential oils from indigenous plants against *Eimeria tenella*. *Vet Parasitol*. 2016 Sep; 228: 96-102. DOI: [10.1016/j.vetpar.2016.08.020](https://doi.org/10.1016/j.vetpar.2016.08.020)
42. Salehi B, Mishra AP, Shukla I, Sharifi-Rad M, Contreras MdM, Segura-Carretero A, et al. Thymol, thyme, and other plant sources: Health and potential uses. *Phytother Res*. 2018 Sep; 32(9): 1688-1706. DOI: [10.1002/ptr.6109](https://doi.org/10.1002/ptr.6109)
43. El-Shall NA, Abd El-Hack ME, Albaqami NM, Khafaga AF, Taha AE, Swelum AA, et al. Phytochemical control of poultry coccidiosis: A review. *Poult Sci*. 2022 Jan; 101(1): 101542. DOI: [10.1016/j.psc.2021.101542](https://doi.org/10.1016/j.psc.2021.101542)

- 10.1016/j.psj.2021.101542
44. Arafa WM, Abolhadid SM, Moawad A, Abdelaty AS, Moawad UK, Shokier KA, et al. Thymol efficacy against coccidiosis in pigeon (*Columba livia domestica*). Prev Vet Med. 2020 Mar; 176: 104914. DOI: 10.1016/j.prevetmed.2020.104914
45. Gholami-Ahangaran M, Ahmadi-Dastgerdi A, Azizi S, Basiratpour A, Zokaei M, and Derakhshan M. Thymol and carvacrol supplementation in poultry health and performance. Vet Med Sci. 2022 Jan; 8(1): 267-288. DOI: 10.1002/vms.3.663
46. Abu El Ezz N, Aboelsoued D, Hassan S, Abdel Megeed K, and El-Metenawy T. Therapeutic effect of *Moringa oleifera* and *Thymus vulgaris* oils against hepatic coccidiosis in experimentally infected rabbits. Trop Biomed. 2020 Dec; 37(4): 1018-1028. DOI: 10.47665/tb.37.4.1018
47. Rubió L, Motilva MJ, and Romero MP. Recent advances in biologically active compounds in herbs and spices: A review of the most effective antioxidant and anti-inflammatory active principles. Crit Rev Food Sci Nut. 2013; 53(9): 943-953. DOI: 10.1080/10408398.2011.574802
48. Alloui MN, Agabou A, and Alloui N. Application of herbs and phytochemical feed additives in poultry production-A Review. Glob J Anim Sci Res. 2014 Jun; 2(3): 234-243. Available at: <http://archives.gjasr.com/index.php/GJASR/article/view/57/155>
49. Peek H, and Landman W. Coccidiosis in poultry: Anticoccidial products, vaccines and other prevention strategies. Vet Q 2011 Sep; 31(3): 143-161. DOI: 10.1080/01652176.2011.605247
50. Mohiti-Asli M, and Ghanaatparast-Rashti M. Dietary oregano essential oil alleviates experimentally induced coccidiosis in broilers. Prev Vet Med. 2015 Jun; 120(2): 195-202. DOI: 10.1016/j.prevetmed.2015.03.014
51. Gordillo Jaramillo FX, Kim DH, Lee SH, Kwon SK, Jha R, and Lee KW. Role of oregano and Citrus species-based essential oil preparation for the control of coccidiosis in broiler chickens. J Anim Sci Biotechnol. 2021 Apr; 12(1): 47. DOI: 10.1186/s40104-021-00569-z
52. Tsinas A, Giannenas I, Voidarou C, Tzora A, and Skoufos J. Effects of an oregano based dietary supplement on performance of broiler chickens experimentally infected with *Eimeria acervulina* and *Eimeria maxima*. J Poultry Sci. 2011 Apr; 48(3): 194-200. DOI: 10.2141/jpsa.010123
53. Alp M, Midilli M, Kocabağlı N, Yılmaz H, Turan N, Gargılı A, et al. The effects of dietary oregano essential oil on live performance, carcass yield, serum immunoglobulin G level, and oocyst count in broilers. J Appl Poult Res. 2012 Sep; 21(3): 630-636. DOI: 10.3382/japr.2012-00551
54. Betancourt L, Hume M, Rodríguez F, Nisbet D, Sohail MU, and Afanador-Tellez G. Effects of Colombian oregano essential oil (*Lippia origanoides* Kunth) and *Eimeria* species on broiler production and cecal microbiota. Poult Sci. 2019 Oct; 98(10): 4777-4786. DOI: 10.3382/ps/pez193
55. Yu M, Jeon JO, Cho HM, Hong JS, Kim YB, Nawarathne SR, et al. Broiler responses to dietary 3, 4, 5-trihydroxybenzoic acid and oregano extracts under *Eimeria* challenge conditions. J Anim Sci Technol. 2021 Nov; 63(6): 1362. DOI: 10.5187%2Fjast.2021.e121
56. Giannenas I, Florou-Paneri P, Papazahariadou M, Botsoglou N, Christaki E, and Spais A. Effect of diet supplementation with ground oregano on performance of broiler chickens challenged with *Eimeria tenella*. Arch Geflügelk. 2004 Jul; 68(6): 247-252. Available at: <https://go.rovedar.com/368>
57. Abbas RZ, Colwell DD, Gilleard J. Botanicals: an alternative approach for the control of avian coccidiosis. World's Poult Sci J. 2012 Jun; 68(2): 203-215. DOI: 10.1017/S0043933912000268
58. Barbour E, Ayyash D, Iyer A, Harakeh S, and Kumosani T. A review of approaches targeting the replacement of coccidiostat application in poultry production. Braz J Poultry Sci. 2015 Oct-Dec; 17: 405-418. DOI: 10.1590/1516-635x1704405-418
59. Shihab IM, AL-Zuhariy M, Abdullah SM, and Mutar SS. Impact of supplementation Neem powder (*Azadirachta indica*) to diet broiler in immunological, physiological and productive traits. Adv Environ Biol. 2017 Mar; 11(3): 44-451.
60. Ishaq R, Chand N, Khan RU, Saeed M, Laudadio V, and Tufarelli V. Methanolic extract of neem (*Azadirachta indica*) leaves mitigates experimentally induced coccidiosis challenge in Japanese quails. J Appl Anim Res. 2022 Aug; 50(1): 498-503. DOI: 10.1080/09712119.2022.2096037
61. Onyiche TE, Gotep JG, Tanko JT, Ochigbo GO, Ozoani HA, Vioff VZ, et al. *Azadirachta indica* aqueous leaf extracts ameliorates coccidiosis in broiler chickens experimentally infected with *Eimeria* oocysts. Sci African. 2021 Jul; 13: e00851.
62. Gotep J, Tanko J, Forcados G, Muraina I, Ozele N, Dogonyaro B, et al. Therapeutic and safety evaluation of combined aqueous extracts of *Azadirachta indica* and *Khaya senegalensis* in chickens experimentally infected with *Eimeria* oocysts. J Parasitol Res. 2016 Feb; 2016:4692424. DOI: 10.1155%2F2016%2F4692424
63. Tipu MA, Pasha T, and Ali Z. Comparative efficacy of salinomycin sodium and neem fruit (*Azadirachta indica*) as feed additive anticoccidials in broilers. Int J Poult Sci. 2002; 1(4): 91-93. DOI: 10.3923/ijps.2002.91.93
64. Dkhil MA, Al-Quraishy S, Abdel Moneim AE, and Delic D. Protective effect of *Azadirachta indica* extract against *Eimeria papillata*-induced coccidiosis. Parasitol Res. 2013 Jan; 112: 101-106. DOI: 10.1007/s00436-012-3109-1
65. Roy R, Hasan M, Aziz F, Islam R, and Sarkar S. Comparative study of neem leaf (*Azadirachta indica*) suspension and toltrazuril against coccidiosis in Sonali chicken. Bangladesh J Vet Med. 2019; 17(2): 97-105. DOI: 10.33109/bjvmd19am1
66. Biu A, Yusuf S, Rabo J. Use of neem (*Azadirachta indica*) aqueous extract as a treatment for poultry coccidiosis in Borno State, Nigeria. Afr Sci. 2021 Sep; 7(3): 147-153. Available at: <https://publications.africanscientistjournal.org/sites/default/files/2022-11/1289-4831-1-PB.pdf>
67. Ramadan R, Youssef F, Mohamed G, Ismail S, El-Bahy M, Abdel-Radi S. Synthesis, characterization and investigation of the anti-coccidial activity of new formulation curcumin-olive oil nano-composite. Adv Anim Vet Sci. 2022 Oct; 10(10): 2186-2196. DOI: 10.17582/journal.aavs/2022/10.8.1781.1786
68. Tajodini M, Saeedi H, and Moghbeli P. Use of black pepper, cinnamon and turmeric as feed additives in the poultry industry. World's Poult Sci J. 2015; 71(1): 175-183. DOI: 10.1017/S0043933915000148
69. Yadav S, Teng PY, Dos Santos TS, Gould RL, Craig SW, Fuller AL, et al. The effects of different doses of curcumin compound on growth performance, antioxidant status, and gut health of broiler chickens challenged with *Eimeria* species. Poult Sci. 2020 Nov; 99(11): 5936-5945. DOI: 10.1016/j.psj.2020.08.046
70. Rajput N, Ali S, Naem M, Khan M, and Wang T. The effect of dietary supplementation with the natural carotenoids curcumin and lutein on pigmentation, oxidative stability and quality of meat from broiler chickens affected by a coccidiosis challenge. Br Poult Sci. 2014 Aug; 55(4): 501-509. DOI: 10.1080/00071668.2014.925537
71. Gogoi C, Sarma J, Barua C, Tamuly S, Upadhyaya T, Islam S, et al. Evaluation of nano-curcumin on experimentally induced coccidiosis in broiler chicks. Int J Chem Stud. 2019 Apr; 7(3): 4514-4520. Available at: <https://www.chemjournal.com/archives/2019/vol7issue3/PartBV/7-3-496-319.pdf>
72. Galli GM, Gerbet RR, Griss LG, Fortuoso BF, Petrolini TG, Boiago MM, et al. Combination of herbal components (curcumin, carvacrol, thymol, cinnamaldehyde) in broiler chicken feed: Impacts on response parameters, performance, fatty acid profiles, meat quality and control of coccidia and bacteria. Microb Pathog. 2020 Feb; 139: 103916. DOI: 10.1016/j.micpath.2019.103916
73. Fatemi A, Asasi K, and Razavi SM. Anticoccidial effects of *Artemisia annua* ethanolic extract: Prevention, simultaneous challenge-medication, and treatment. Parasitol Res. 2017 Sep; 116: 2581-2589. DOI: 10.1007/s00436-017-5567-y
74. Jiao J, Yang Y, Liu M, Li J, Cui Y, Yin S, et al. Artemisinin and *Artemisia annua* leaves alleviate *Eimeria tenella* infection by facilitating apoptosis of host cells and suppressing inflammatory response. Vet Parasitol. 2018 Apr; 254: 172-177. DOI: 10.1016/j.vetpar.2018.03.017
75. Fatemi A, Razavi SM, Asasi K, and Torabi Goudarzi M. Effects of *Artemisia annua* extracts on sporulation of *Eimeria* oocysts. Parasitol Res. 2015 Mar; 114: 1207-1211. DOI: 10.1007/s00436-014-4304-z
76. Azeez TB, and Lunghar J. Antiinflammatory effects of turmeric (*Curcuma longa*) and ginger (*Zingiber officinale*). Inflamm Natu Prod. 2021; 127-146. DOI: 10.1016/B978-0-12-819218-4.00011-0
77. Lahlou RA, Bounechada M, Mohammedi A, Silva LR, and Alves G. Dietary use of *Rosmarinus officinalis* and *Thymus vulgaris* as anticoccidial alternatives in poultry. Anim Feed Sci Technol. 2021 Jan; 273: 114826. Available at: <https://www.x-mol.net/paper/article/1349111312502517760>
78. Islas JF, Acosta E, Zuca G, Delgado-Gallegos JL, Moreno-Treviño MG,

- Escalante B, et al .An overview of Neem (*Azadirachta indica*) and its potential impact on health. J Funct Food. 2020 Nov; 74: 104171. DOI: [10.1016/j.jff.2020.104171](https://doi.org/10.1016/j.jff.2020.104171)
79. Sarkar S, Singh RP, and Bhattacharya G. Exploring the role of *Azadirachta indica* (neem) and its active compounds in the regulation of biological pathways: An update on molecular approach. 3 Biotech. 2021 Apr; 11(4): 178. DOI: [10.1007/s13205-021-02745-4](https://doi.org/10.1007/s13205-021-02745-4)
80. Calixto JB. The role of natural products in modern drug discovery. An Acad Bras Ciênc. 2019; 91(suppl3): e20190105. DOI: [10.1590/0001-3765201920190105](https://doi.org/10.1590/0001-3765201920190105)
81. Heinrich M, Appendino G, Efferth T, Fürst R, Izzo AA, Kayser O, et al. Best practice in research—overcoming common challenges in phytopharmacological research. J Ethnopharmacol. 2020 Jan; 246: 112230. DOI: [10.1016/j.jep.2019.112230](https://doi.org/10.1016/j.jep.2019.112230)
82. Sen S, and Chakraborty R. Revival, modernization and integration of Indian traditional herbal medicine in clinical practice: Importance, challenges and future. J Tradit Complement Med. 2017 Jun; 7(2): 234-244. DOI: <https://doi.org/10.1016/j.jtcme.2016.05.006>
83. Jamshidi-Kia F, Lorigooini Z, and Amini-Khoei H. Medicinal plants: Past history and future perspective. J HerbMed Pharmacol. 2017; 7(1): 1-7. DOI: [10.15171/jhp.2018.01](https://doi.org/10.15171/jhp.2018.01)
84. Atanasov AG, Waltenberger B, Pferschy-Wenzig EM, Linder T, Wawrosch C, Uhrin P, et al. Discovery and resupply of pharmacologically active plant-derived natural products: A review. Biotechnol Adv. 2015 Dec; 33(8): 1582-1614. DOI: [10.1016/j.biotechadv.2015.08.001](https://doi.org/10.1016/j.biotechadv.2015.08.001)
85. David B, Wolfender JL, and Dias DA. The pharmaceutical industry and natural products: Historical status and new trends. Phytochem Rev. 2015 Jun; 14: 299-315. Available at: <https://link.springer.com/article/10.1007/s11101-014-9367-z>
86. Hafez HM, and Attia YA. Challenges to the poultry industry: Current perspectives and strategic future after the COVID-19 outbreak. Front Vet Sci. 2020 Aug; 7: 516. DOI: [10.3389%2Ffvets.2020.00516](https://doi.org/10.3389%2Ffvets.2020.00516)
87. Withers PJ, Elser JJ, Hilton J, Ohtake H, Schipper WJ, and Van Dijk KC. Greening the global phosphorus cycle: How green chemistry can help achieve planetary P sustainability. Green Chem. 2015 Mar; 17(4): 2087-2099. Available at: <https://pubs.rsc.org/en/content/articlelanding/2015/gc/c4gc02445a>